

Dictation Note (Blank Template)

January 1, 2050

Name: Joe Sample

Record #: 2813308004

Date:

Pre-Op Diagnosis:

Post-Op Diagnosis:

Procedure:

Surgeon: Dr. Mallet DPM

Assistant:

Anesthesia:

Hemostasis:

Report:

Before surgery

“The patient was brought into the OR suite and placed in the (supine, prone, lateral) position. (MAC, general, spinal) anesthesia was appropriately induced. A pneumatic (ankle/thigh) tourniquet was placed on the (left/right) extremity. Our attention was then directed to the (left/right) site which we noted a (skin tenting, abscess, bunion, deformity, fracture, etc.). Using 20cc of 1:1 mixture of 1% Lidocaine plain and _____cc of 0.5% Marcaine plain, we performed a (Local, Common Peroneal, PT) block in the (MAYO, Ankle block, isolated) fashion. The foot was scrubbed, prepped, and draped in an aseptic manner. An Esmarch bandage was used to exsanguinate

the (left/right) foot. The (ankle/thigh) tourniquet was inflated and set to _____ mmHg.”

During Surgery

“At this time, attention was directed to the dorsal aspect of the first metatarsal head of the (left/right) foot where a 5cm linear longitudinal incision was made medial to the EHL, following the contour of the deformity. The incision was deepened using blunt dissection. All vital neural and vascular structures were identified. All bleeders were cauterized with Bovie as necessary...”

After Surgery

“The tourniquet was deflated, and it was noted a hyperemic response of the (left/right) lower extremity. A postoperative block consisted of _____ cc’s of 0.5% Marcaine plain. The incision site was then dressed in _____ (Betadine soaked adaptic, Xeroform, gauze, Kling, Kerlix, steri-strips). A posterior splint, post-op shoe, CAM boot were then applied. The patient will be discharged home with written instructions.”